

Metabolic tracing unravels pathways of fungal and bacterial amino sugar formation in soil

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 British Society of Soil Science The present study aimed to reveal the steps in the formation of fungal and bacterial amino sugars (AS) from glucose. Glucose labelled uniformly and position-specifically at C-1, C-2, C-4 and C-6 was applied to an agricultural soil. Fungal and bacterial pathways of AS formation were reconstructed by a new approach: ^{13}C recovery from individual C positions of glucose in AS by IC-O-IRMS. Only 0.75% of ^{13}C from initially applied glucose was incorporated into AS within 10 days and followed the order: glucosamine > galactosamine > muramic acid. Relative incorporation of ^{13}C (% of AS-C) had the largest values for bacterial muramic acid and the smallest for fungal galactosamine. This reflects faster turnover of bacterial than fungal cell walls. A maximum of $55\% \pm 19\%$ of the ^{13}C incorporated in glucosamine was derived from intact, untransformed glucose. The smallest recovery was observed for C-1 and C-4. Recovery of C-1 even decreased from day 3 to day 10, reflecting the metabolization by the pentose phosphate pathway. To be incorporated into AS, metabolites resulting from either the pentose phosphate pathway or glycolysis needed to be reconstructed to glucose by the reversible pathways of gluconeogenesis. The direct formation of AS (e.g. intact recovery of the glucose precursor), as well as amino sugar formation from recycled metabolites allocated to the hexose pool by back flux, occurred in parallel. Bacterial muramic acid showed the most variation in the recovery of individual C positions during 10 days, which reflects its intensive transformation by glycolysis, pentose phosphate pathway and gluconeogenesis. It is explained by the lower metabolic activity of fungi than bacteria at steady-state conditions. The present approach enabled us to analyse the biochemical pathways in bacteria and fungi, which will considerably improve our understanding of the formation and stabilization of microbially derived SOM compounds. Highlights: Fungal and bacterial pathways of the formation of amino sugars from glucose are reconstructed Intact use of glucose is distinguished from recycling by a metabolic tracing approach Half of the precursor (glucose) is metabolized prior to incorporation into amino sugars Intracellular recycling plays a crucial role in microbial SOM formation.

<http://dx.doi.org/10.1111/ejss.12736>

References

- [1] Amelung, W. 2003. Nitrogen biomarkers and their fate in soil. *Journal of Plant Nutrition and Soil Science*, 166, 677–686.
- [2] Apostel, C., Dippold, M., Glaser, B. & Kuzyakov, Y. 2013. Biochemical pathways of amino acids in soil: assessment by position-specific labeling and C-PLFA analysis. *Soil Biology & Biochemistry*, 67, 31–40.
- [3] Apostel, C., Dippold, M.A. & Kuzyakov, Y. 2015. Biochemistry of hexose and pentose transformations in soil analyzed by position-specific labeling and C-PLFA. *Soil Biology & Biochemistry*, 80, 199–208.

- [4] Beckham, G.T., Matthews, J.F., Peters, B., Bomble, Y.J., Himmel, M.E. & Crowley, M.F. 2011. Molecular-level origins of biomass recalcitrance: decrystallization free energies for four common cellulose polymorphs. *Journal of Physical Chemistry*, 115, 4118-4127.
- [5] Bode, S., Denef, K. & Boeckx, P. 2009. Development and evaluation of a high-performance liquid chromatography/isotope ratio mass spectrometry methodology for δC analyses of amino sugars in soil. *Rapid Communications in Mass Spectrometry*, 23, 2519-2526.
- [6] Bode, S., Fancy, R. & Boeckx, P. 2013. Stable isotope probing of amino sugars - a promising tool to assess microbial interactions in soils. *Rapid Communications in Mass Spectrometry*, 27, 1367-1379.
- [7] Cadisch, G. & Giller, K.E. 1996. Estimating the contribution of legumes to soil organic matter build up in mixed communities of C3/C4 plants. *Soil Biology & Biochemistry*, 28, 823-825.
- [8] Caspi, R., Foerster, H., Fulcher, C.A., Kaipa, P., Krummenacker, M., Latendresse, M. et al. 2008. The MetaCyc database of metabolic pathways and enzymes and the BioCyc collection of pathway/genome databases. *Nucleic Acids Research*, 36, 623-631.
- [9] Derrien, D., Marol, C. & Balesdent, J. 2007. Microbial biosyntheses of individual neutral sugars among sets of substrates and soils. *Geoderma*, 139, 190-198.
- [10] Dijkstra, P., Blankinship, J.C., Selmants, P.C., Hart, S.C., Koch, G.W., Schwartz, E. et al. 2011. Probing carbon flux patterns through soil microbial metabolic networks using parallel position-specific tracer labeling. *Soil Biology & Biochemistry*, 43, 126-132.
- [11] Dippold, M.A. & Kuzyakov, Y. 2016. Direct incorporation of fatty acids into microbial phospholipids in soils: position-specific labeling tells the story. *Geochimica et Cosmochimica Acta*, 174, 211-221.
- [12] Dippold, M.A., Boesel, S., Gunina, A., Kuzyakov, Y. & Glaser, B. 2014. Improved δC -analysis of amino sugars in soil by ion chromatography-oxidation-isotope ratio mass spectrometry. *Rapid Communications in Mass Spectrometry*, 28, 569-576.
- [13] Engelking, B., Flessa, H. & Joergensen, R.G. 2007. Shifts in amino sugar and ergosterol contents after addition of sucrose and cellulose to soil. *Soil Biology & Biochemistry*, 39, 2111-2118.
- [14] Fernandez, C.W. & Koide, R.T. 2012. The role of chitin in the decomposition of ectomycorrhizal fungal litter. *Ecology*, 93, 24-28.
- [15] Fokin, A.D., Knyazev, D.A. & Kuzyakov, Y.V. 1993. Destruction of C-labeled and N-labeled amino-acids and nucleic bases in soil and the supply of their transformation products to plants. *Eurasian Soil Science*, 25, 109-122.
- [16] Gearing, P.J., Gearing, J.N., Maughan, J.T. & Oviatt, C.A. 1991. Isotopic distribution of carbon from sewage-sludge and eutrophication in the sediments and food web of estuarine ecosystems. *Environmental Science & Technology*, 25, 295-301.
- [17] Glaser, B. & Amelung, W. 2002. Determination of C-natural abundance of amino acid enantiomers in soil: methodological considerations and first results. *Rapid Communications in Mass Spectrometry*, 16, 891-898.
- [18] Glaser, B. & Gross, S. 2005. Compound-specific δC -analysis of individual amino sugars - a tool to quantify timing and amount of soil microbial residue stabilization. *Rapid Communications in Mass Spectrometry*, 19, 1409-1416.
- [19] Glaser, B., Turrion, M.B. & Alef, K. 2004. Amino sugars and muramic acid - biomarkers for soil microbial community structure analysis. *Soil Biology & Biochemistry*, 36, 399-407.
- [20] Glaser, B., Millar, N. & Blum, H. 2006. Sequestration and turnover of bacterial- and fungal-derived carbon in a temperate grassland soil under long-term elevated atmospheric pCO₂. *Global Change Biology*, 12, 1521-1531.
- [21] Gottwald, W. 2000. Statistik für Anwender. 1. Auflage. Wiley-VCH-Verlag, Weinheim.
- [22] Gunina, A., Smith, A.R., Kuzyakov, Y. & Jones, D.L. 2017a. Microbial uptake and utilization of low molecular weight organic substrates in soil depend on carbon oxidation state. *Biogeochemistry*, 133, 89-100.
- [23] Gunina, A., Dippold, M., Glaser, B. & Kuzyakov, Y. 2017b. Turnover of microbial groups and cell components in soil: C-analysis of cellular biomarkers. *Biogeosciences*, 14, 271-283.
- [24] Indorf, C., Bode, S., Boeckx, P., Dyckmans, J., Meyer, A., Fischer, K. et al. 2013. Comparison of HPLC methods for the determination of amino sugars in soil hydrolysates. *Analytical Letters*, 46, 2145-2164.
- [25] Keiluweit, M., Bougoure, J.J., Zeglin, L.H., Myrold, D.D., Weber, P.K., Pett-Ridge, J. et al. 2013. Nano-scale investigation of the association of microbial nitrogen residues with iron hydroxides in a forest soil O-horizon. *Geochimica et Cosmochimica Acta*, 95, 213-226.
- [26] Keseler, I.M., Bonavides-Martinez, C., Collado-Vides, J., Gama-Castro, S., Gunsalus, R.P., Johnson, D.A. et al. 2009. EcoCyc: a comprehensive view of Escherichia coli biology. *Nucleic Acids Research*, 37, 464-470.
- [27] Kotnik, M., Anderluh, P.S. & Prezelj, A. 2007. Development of novel inhibitors targeting intracellular steps of peptidoglycan biosynthesis. *Current Pharmaceutical Design*, 13, 2283-2309.
- [28] Kuzyakov, Y. & Domanski, G. 2000. Carbon input by plants into the soil. Review. *Journal of Plant Nutrition and Soil Science*, 163, 421-431.
- [29] Lengeler, J.W., Drews, G. & Schlegel, H.G. 1999. Biology of the Prokaryotes. Georg Thieme Verlag, Stuttgart.

- [30] von Luetzow, M., Kögel-Knabner, I., Ekschmitt, K., Matzner, E., Guggenberger, G., Marschner, B. et al. 2006. Stabilization of organic matter in temperate soils: mechanisms and their relevance under different soil conditions—a review. *European Journal of Soil Science*, 57, 426–445.
- [31] Macura, J. & Kubatova, Z. 1973. Control of carbohydrate utilization by soil microflora. *Soil Biology & Biochemistry*, 5, 193–204.
- [32] Merzendorfer, H. 2011. The cellular basis of chitin synthesis in fungi and insects: common principles and differences. *European Journal of Cell Biology*, 90, 759–769.
- [33] Meyer, A., Fischer, H., Kuzyakov, Y. 2008. Improved RP-HPLC and anion-exchange chromatography methods for the determination of amino acids and carbohydrates in soil solutions. *Journal of Plant Nutrition & Soil Science (Zeitschrift Fur Pflanzenernahrung Und Bodenkunde)*, 171, 917–926.
- [34] Milewski, S., Gabriel, L. & Olchow, J. 2006. Enzymes of UDP-GlcNAc biosynthesis in yeast. *Yeast*, 23, 1–14.
- [35] Miltner, A., Bombach, B., Schmidt-Brücken, B. & Kästner, M. 2012. SOM genesis: microbial biomass as a significant source. *Biogeochemistry*, 111, 41–55.
- [36] Moore, J.C., McCann, K. & de Ruiter, P.C. 2005. Modeling trophic pathways, nutrient cycling, and dynamic stability in soils. *Pedobiologia*, 49, 499–510.
- [37] Park, J.T. & Uehara, T. 2008. How bacteria consume their own exoskeletons (turnover and recycling of cell wall peptidoglycan). *Microbiology and Molecular Biology, Reviews*, 72, 211–227.
- [38] Rannekleiv, S.B. & Baath, E. 2003. Use of phospholipid fatty acids to detect previous self-heating events in stored peat. *Applied and Environmental Microbiology*, 69, 3532–3539.
- [39] Rethemeyer, J., Kramer, C., Gleixner, G., Wiesenberger, G.L.B., Schwark, L., Andersen, N. et al. 2004. Complexity of soil organic matter: AMS C- analysis of soil lipid fractions and individual compounds. *Radiocarbon*, 46, 465–473.
- [40] Richmond, M.H. & Perkins, H.R. 1962. Biosynthesis of muramic acid by intact staphylococci. *Biochemical Journal*, 85, 580–588.
- [41] Scandellari, F., Hobbie, E.A., Ouimette, A.P. & Stucker, V.K. 2009. Tracing metabolic pathways of lipid biosynthesis in ectomycorrhizal fungi from position-specific C-labelling in glucose. *Environmental Microbiology*, 11, 3087–3095.
- [42] Uehara, T. & Park, J.T. 2008. Growth of *Escherichia coli*: significance of peptidoglycan degradation during elongation and septation. *Journal of Bacteriology*, 190, 3914–3922.
- [43] Veuger, B., van Oevelen, D., Boschker, H.T.S. & Middelburg, J.J. 2006. Fate of peptidoglycan in an intertidal sediment: an in situ C-labeling study. *Limnology and Oceanography*, 51, 1572–1580.
- [44] Waring, B.G., Averill, C. & Hawkes, C.V. 2013. Differences in fungal and bacterial physiology alter soil carbon and nitrogen cycling: insights from meta-analysis and theoretical models. *Ecology Letters*, 16, 887–894.